

FUNDAMENTALS OF LOGIC

NO.5 SOUNDNESS AND COMPLETENESS

Tatsuya Hagino

hagino@sfc.keio.ac.jp

lecture URL

<https://vu5.sfc.keio.ac.jp/slide/>

So Far

- Propositional Logic
 - Logical Connectives (\wedge , \vee , \rightarrow , \neg)
 - Truth Table
 - Tautology
- Normal Form
 - Disjunctive Normal Form
 - Conjunctive Normal Form
 - Restricting Logical Connectives
- Proof
 - Axiom and Theorem
 - LK Framework

Tautology for Sequent

- Extend the notion of tautology to **sequent**.
- Let Γ be a sequence of formulae A_1, \dots, A_m :

$$\Gamma^* = \begin{cases} A_1 \vee \dots \vee A_m & \text{when } m > 0 \\ \perp & \text{when } m = 0 \end{cases}$$

$$\Gamma_* = \begin{cases} A_1 \wedge \dots \wedge A_m & \text{when } m > 0 \\ \top & \text{when } m = 0 \end{cases}$$

- Sequent $\Gamma \vdash \Delta$ is a **tautology** $\Leftrightarrow \Gamma_* \rightarrow \Delta^*$ is a tautology.

Soundness and Completeness

- $\Gamma \vdash \Delta$ is a **tautology**:
 - For any assignment v , $v(\Gamma_* \rightarrow \Delta^*) = T$
 - For any assignment of truth values to propositional variables, we can
 - calculate the truth value of $\Gamma_* \rightarrow \Delta^*$ using the truth tables and check whether it is always true.
- $\Gamma \vdash \Delta$ is **provable** in LK:
 - There is a proof figure of which end sequent is $\Gamma \vdash \Delta$.
 - Starting from initial sequents, we can construct a proof figure of $\Gamma \vdash \Delta$.
- **Soundness**
 - Any sequent which is provable in LK is a tautology.
- **Completeness**
 - Any tautology is provable in LK.

LK Soundness Theorem

Theorem: For any sequent $\Gamma \vdash \Delta$, if $\Gamma \vdash \Delta$ is provable in LK, $\Gamma \vdash \Delta$ is a tautology.

Proof: We need to show the following two things:

1. An initial sequent is a tautology.
2. For each inference rule, if all the sequents above are tautologies, the bottom sequent is also a tautology.

For (1), an initial sequent $A \vdash A$ is $A \rightarrow A$, and it is a tautology.

For (2), we need to check for each inference rule. For example,

$$\frac{\Gamma_1 \vdash \Delta_1, A \quad B, \Gamma_2 \vdash \Delta_2}{A \rightarrow B, \Gamma_1, \Gamma_2 \vdash \Delta_1, \Delta_2} (\rightarrow L)$$

we need to show when $\Gamma_{1*} \rightarrow \Delta_{1*} \vee A$ and $B \wedge \Gamma_{2*} \rightarrow \Delta_{2*}$ are tautologies, $((A \rightarrow B) \wedge \Gamma_{1*} \wedge \Gamma_{2*}) \rightarrow (\Delta_{1*} \vee \Delta_{2*})$ is also a tautology.

cont.

For an assignment v if $v((A \rightarrow B) \wedge \Gamma_{1*} \wedge \Gamma_{2*}) = T$, then $v(A \rightarrow B) = T$, $v(\Gamma_{1*}) = T$, and $v(\Gamma_{2*}) = T$.

1. If $v(A) = F$, from $v(\Gamma_{1*} \rightarrow \Delta_1^* \vee A) = T$ and $v(\Gamma_{1*}) = T$, $v(\Delta_1^*) = T$.
2. If $v(A) = T$, from $v(A \rightarrow B) = T$, $v(B) = T$. Since $v(B \wedge \Gamma_{2*} \rightarrow \Delta_2^*) = T$ and $v(\Gamma_{2*}) = T$, $v(\Delta_2^*) = T$.

In both cases, $v(\Delta_1^* \vee \Delta_2^*) = T$. Therefore, $((A \rightarrow B) \wedge \Gamma_{1*} \wedge \Gamma_{2*}) \rightarrow (\Delta_1^* \vee \Delta_2^*)$ is a tautology


For other rules, we can show the bottom sequent is a tautology when the above sequents are tautologies.

Therefore LK framework is sound. QED

LK Completeness Theorem

Theorem: If $\Gamma \vdash \Delta$ is a tautology, $\Gamma \vdash \Delta$ is provable in LK without using cut inference rule.

Proof: First, define the **decomposition** of a sequent as follows:




A sequent is decomposed into one or two sequents.

Two things hold for any decomposition:

- The number of logical connectives decreases.
- If the original sequent is a tautology, the decompositions are tautologies.

cont.

Apply decomposition repeatedly:



Sequents at the bottom row do not contain logical connectives.

- **Complete decomposition tree**
- If $\Gamma \vdash \Delta$ is a tautology, the bottom row of the complete decomposition contains tautology sequents containing propositional variables (no logical connectives).

A sequent $p_1, \dots, p_m \vdash q_1, \dots, q_n$ containing only propositional variables is a tautology if and only if one of p_i matches with one of q_j .


$\Gamma_1, A, \Gamma_2 \vdash \Delta_1, A, \Delta_2$ is provable without cut rule.

- Apply weakening and exchange rules to initial sequent $A \vdash A$.

cont.

When $\Gamma \vdash \Delta$ is decomposed into $\Gamma_1 \vdash \Delta_1$ and $\Gamma_2 \vdash \Delta_2$, if $\Gamma_1 \vdash \Delta_1$ and $\Gamma_2 \vdash \Delta_2$ are provable without cut, $\Gamma \vdash \Delta$ is also provable without cut.

- For each decomposition, use rules for logical connectives, exchange and contraction.



Therefore, if a sequent is a tautology, the bottom row sequents of its complete decomposition are tautologies and they are provable without cut. By traversing back the decomposition, we can see that the original sequent is also provable without cut.

- On the other hand, if a given sequent is not a tautology, no matter how we may decompose it, there exists a non tautology sequent at the bottom row.

LK framework is complete. QED

This proof provides an algorithm to check whether a sequent is provable or not.

Example

- Find a complete decomposition of $\vdash ((p \rightarrow q) \rightarrow p) \rightarrow p$ and create a proof figure without cut.

Decomposition

$\vdash ((p \rightarrow q) \rightarrow p) \rightarrow p$



Proof

$\vdash ((p \rightarrow q) \rightarrow p) \rightarrow p$

Cut Elimination Theorem

Theorem: If $\Gamma \vdash \Delta$ is provable in LK, there exists a proof figure of $\Gamma \vdash \Delta$ without cut.

Proof: Since any provable sequent is a tautology, it is provable without cut. QED

A proof figure with cut

$$\begin{array}{c}
 \frac{\frac{\frac{\frac{\frac{A \vdash A}{\vdash A, \neg A}}{\vdash A, A \vee \neg A}}{\vdash A \vee \neg A, A}}{\vdash A \vee \neg A, A \vee \neg A}}{\vdash A \vee \neg A} \\
 \frac{\frac{\frac{\frac{A \vdash A}{B, A \vdash A}}{A \vdash B \rightarrow A}}{A \vdash (A \rightarrow B) \vee (B \rightarrow A)} \\
 \frac{\frac{\frac{\frac{A \vdash A}{\neg A, A \vdash}}{A, \neg A \vdash}}{A, \neg A \vdash B}}{\neg A \vdash A \rightarrow B}}{\neg A \vdash (A \rightarrow B) \vee (B \rightarrow A)} \\
 \hline
 \frac{A \vee \neg A \vdash (A \rightarrow B) \vee (B \rightarrow A)}{\vdash (A \rightarrow B) \vee (B \rightarrow A)}
 \end{array}$$

Theorem: Any formula in a cut free proof figure of $\Gamma \vdash \Delta$ is a sub formula of $\Gamma \vdash \Delta$.

Exercise

- Find a proof figure of $\vdash (A \rightarrow B) \vee (B \rightarrow A)$ without cut.

Duality

- Duality in inference rules:
 - $(\neg L)$ and $(\neg R)$ is symmetrical by exchanging left and right of $\Gamma \vdash \Delta$.

$$\frac{\Gamma \vdash \Delta, A}{\neg A, \Gamma \vdash \Delta} (\neg L)$$

$$\frac{A, \Gamma \vdash \Delta}{\Gamma \vdash \Delta, \neg A} (\neg R)$$

- Inference rules for \vee and \wedge are symmetrical by replacing left and right and \vee and \wedge .

$$\frac{A, \Gamma \vdash \Delta}{A \wedge B, \Gamma \vdash \Delta} (\wedge L_1)$$

$$\frac{\Gamma \vdash \Delta, A}{\Gamma \vdash \Delta, A \vee B} (\vee R_1)$$

$$\frac{B, \Gamma \vdash \Delta}{A \wedge B, \Gamma \vdash \Delta} (\wedge L_2)$$

$$\frac{\Gamma \vdash \Delta, B}{\Gamma \vdash \Delta, A \vee B} (\vee R_2)$$

$$\frac{A, \Gamma_1 \vdash \Delta_1 \quad B, \Gamma_2 \vdash \Delta_2}{A \vee B, \Gamma_1, \Gamma_2 \vdash \Delta_1, \Delta_2} (\vee L)$$

$$\frac{\Gamma_1 \vdash \Delta_1, A \quad \Gamma_2 \vdash \Delta_2, B}{\Gamma_1, \Gamma_2 \vdash \Delta_1, \Delta_2, A \wedge B} (\wedge R)$$

Duality Theorem:

Let A and B be formulae without \rightarrow . Let \tilde{A} and \tilde{B} be formulae A and B but replacing \wedge and \vee . If $A \vdash B$ is provable, $\tilde{B} \vdash \tilde{A}$ is also provable.

Summary

- Soundness of LK framework
 - Any sequent which is provable in LK is a tautology.
- Completeness of LK framework
 - A tautology is provable in LK.
 - There is an algorithm of checking whether a formula is provable or not.
- Properties of LK framework
 - cut elimination theorem
 - duality theorem