

# KA-wavelets on semisimple Lie groups and quasi-orthogonality of matrix coefficients

Takeshi Kawazoe  
Keio University at Fujisawa  
5322 Endo, Fujisawa, 252-8520 Japan  
kawazoe@sfc.keio.ac.jp

## §1 Introduction.

First we brief the history of continuous wavelet transforms. Originally the (continuous) wavelet transform, introduced by Morlet around 1980, was the following one. We denote by  $H^2(\mathbf{R})$  the closed subspace of  $L^2(\mathbf{R})$  consisting of all  $L^2$  functions  $f$  on  $\mathbf{R}$  with  $\text{supp}(\hat{f}) \subset [0, \infty)$ , and we fix  $\psi \in H^2(\mathbf{R})$  satisfying the so-called admissible condition

$$c_\psi = \int_0^\infty \frac{|\hat{\psi}(\lambda)|^2}{\lambda} d\lambda < \infty.$$

Then the wavelet transform  $W_\psi$  associated to  $\psi$  is defined on  $H^2(\mathbf{R})$  as

$$W_\psi f(u, v) = \int_{-\infty}^{\infty} f(x) e^{-u/2} \bar{\psi}(e^{-u}x + v) dx \quad (u, v \in \mathbf{R}).$$

**Theorem 1.1.**  $W_\psi$  is an isometric isomorphism from  $H^2(\mathbf{R})$  onto  $L^2(\mathbf{R}^2)$ : For any  $f \in H^2(\mathbf{R})$

$$\|f\|^2 = \frac{1}{c_\psi} \|W_\psi f\|^2.$$

Furthermore, for any  $f \in H^2(\mathbf{R})$  and  $x \in \mathbf{R}$  at which  $f$  is continuous,

$$f(x) = \frac{1}{c_\psi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (W_\psi f)(u, v) e^{-u/2} \bar{\psi}(e^{-u}x + v) du dv.$$

In [GMP] Grossmann-Morlet-Paul pointed out the group-theoretical interpretation of the wavelet transform  $W_\psi$ . Let  $G$  be the affine group  $\mathbf{R}^2$  with multiplication law:

$$(u, v)(u', v') = (u + u', e^{-u'}v + v'),$$

and let  $(T, H^2(\mathbf{R}))$  be an irreducible unitary representation of  $G$  defined by

$$(T(u, v)f)(x) = e^{-u/2} f(e^{-u}x + v) \quad (f \in H^2(\mathbf{R})).$$

In this scheme  $W_\psi$  can be rewritten as

$$W_\psi f(u, v) = \langle f, T(u, v)\psi \rangle,$$

where  $\langle \cdot, \cdot \rangle$  is the inner product of  $H^2(\mathbf{R})$ . Furthermore, since  $dudv$  is a left invariant Haar measure on  $G$ , Theorem 1.1 yields the square-integrability and the orthogonality of the matrix coefficients  $\langle f, T(u, v)\psi \rangle$  of  $T$  on  $G$ . In this sense the theory of the continuous wavelet transform  $W_\psi$  on  $H^2(\mathbf{R})$  is nothing but the one of the square-integrable representation  $(T, H^2(\mathbf{R}))$  of  $G$ .

General theory of square-integrable representations of locally compact groups has been investigated by various mathematicians; Weyl [W] for compact groups, Godement [G] for unimodular locally compact groups, and Duflo-Moore [DM] for general locally compact groups. Explicit theory based on the construction of the square-integrable representations was obtained by Harish-Chandra [HC] for semisimple Lie groups and by Moore-Wolf [MW] for nilpotent groups.

How to extend the theory of square-integrable representations of locally compact groups  $G$ ? One of the ways is to replace the square-integrability on  $G$  by the one on a quotient space  $G/H$  for a closed subgroup  $H$  of  $G$ . More generally, find a representation  $(T, \mathcal{H})$  of  $G$ , a measurable subset  $(S, ds)$  of  $G$ , and  $\psi \in \mathcal{H}$  for which, for any  $f \in \mathcal{H}$

$$(\star) \quad \|f\|^2 = \frac{1}{c_{S, \psi}} \int_S |\langle f, T(s)\psi \rangle|^2 ds.$$

Then, it is easy to see that the transform defined by  $\langle f, T(s)\psi \rangle$  is an isometric isomorphism from  $\mathcal{H}$  onto  $L^2(S, ds)$ , and each  $f \in \mathcal{H}$  has an  $L^2$  decomposition in the weak sense:

$$f = \frac{1}{c_{S, \psi}} \int_S \langle f, T(s)\psi \rangle T(s)\psi ds.$$

For the last decade researches has been done in this scheme and many wavelet transforms has been constructed on locally compact groups, for example, on  $\mathbf{R}_+^* \times SO(n)$  by Murenzi [M], on  $\mathbf{R}_+^* \times SO(1, n)$  by A-J. Unterberger [U], on  $\mathbf{R}_+^* \times SO(1, n) \times \mathbf{R}^{n+1}$  by Bohnke [B], on  $S \times V$ ,  $V$  is a vector space and  $S$  is

a subgroup of  $GL(V)$ , by De Bièvre [DB], on  $SO(2, 1) \times \mathbf{R}^3$  by Ali, Antoine, Gazeau [AAG], on  $\mathbf{R}_+^* \times SO(n) \times H_n$  by Kalisa-Toréssani [KT], Toréssani [T1,2], on  $GL(n, \mathbf{R})$  by Bernier-Taylor [BT], on  $SO(2, 1)$  by Wu-Zhong [WZ], and on Iwasawa  $AN$  groups by Kawazoe [K3] and Liu [L].

In this paper we shall consider the case that  $G$  is a semisimple Lie group and  $S = KA$ , where  $K$  and  $A$  are respectively the maximal compact and abelian subgroups of  $G$ . More precisely, let  $G$  be a semisimple Lie group with finite center and  $G = KAK$  the Cartan decomposition of  $G$ .  $dg$  denotes a Haar measure on  $G$  and  $dg = D(a)dkdak$  the corresponding decomposition of  $dg$ . Then we take  $S = KA$  and  $ds = D(a)dkda$  in the above scheme, and we try to find a representation  $(T, \mathcal{H})$  of  $G$  and  $\psi \in \mathcal{H}$  satisfying  $(\star)$ . Unfortunately, the condition  $(\star)$  is very strong, so I feel that we have no answer for  $T$  and  $\psi$ . Therefore, we shall consider a weak condition; there exist constants  $0 < C_1, C_2 < \infty$  such that

$$(\star\star) \quad C_1 \|f\|^2 \leq \int_S |\langle f, T(s)\psi \rangle|^2 ds \leq C_2 \|f\|^2$$

and we shall obtain a sufficient condition on  $\psi$  for which  $\langle f, T(s)\psi \rangle$  satisfies  $(\star\star)$  (see Theorem 3.1). In §4 we shall treat the case of  $G = SU(1, 1)$  and  $(T_{1/2}, \mathcal{H}_{1/2})$  the limit of the holomorphic discrete series of  $G$ . We note that  $T_{1/2}$  is not square-integrable on  $G$ . Then we shall find a  $\psi \in \mathcal{H}_{1/2}$  satisfying  $(\star\star)$ . Moreover, we shall deduce that, if we ignore a finite dimensional subspace of  $\mathcal{H}_{1/2}$ , then we can find a  $\psi \in \mathcal{H}_{1/2}$  satisfying  $(\star)$  (see Theorem 4.4). In this process we use the facts that some differences of the matrix coefficients of  $T_{1/2}$  are square-integrable on  $\mathbf{R}$  with respect to  $D(a)da$  and moreover, they satisfy a quasi-orthogonality. These facts are summarized in Lemmas 4.1, 4.2, and 4.3.

After the lecture, the author noticed that J.-P. Antoine and P. Vandergheynst [AV1,2] had the same idea and they obtained an example in the case of  $SO(3, 1)$ .

## §2. Notation.

Let  $G$  be a semisimple Lie group with finite center and  $G = KAN$  the Iwasawa decomposition of  $G$ . Let  $\Sigma$  be the set of roots for  $(G, A)$  and  $\Sigma^+$  the one of positive roots corresponding to  $N$ . Let  $A^+$  denote the closed positive Weyl chamber in  $A$  and  $G = KA^+K$  the Cartan decomposition of  $G$ . Let

$dg$  denote a Haar measure on  $G$ , and  $dk$ ,  $da$ , and  $dn$  ones for  $K$ ,  $A$ , and,  $N$  respectively. We normalize  $dk$  as  $\int_K dk = 1$ . According to the Iwasawa and Cartan decompositions of  $G$ , there are decompositions of  $dg$  such that

$$dg = e^{\rho(\log a)} dk dadn = D(a) dk dadk',$$

where  $\rho = \frac{1}{2} \sum_{\alpha \in \Sigma^+} m_\alpha \alpha$  and

$$D(a) = \prod_{\alpha \in \Sigma^+} (\sinh \alpha(\log a))^{m_\alpha},$$

$m_\alpha$  stands for the multiplicity of  $\alpha$ .

### §3. $KA$ -wavelets.

Let  $(T, \mathcal{H})$  be a unitary representation of  $G$  and

$$\mathcal{H} = \bigoplus_{\tau \in \hat{K}} \mathcal{H}_\tau,$$

the  $K$ -type decomposition of  $\mathcal{H}$ . In the following argument we assume that

$$[T, \tau] \leq 1,$$

and we denote by  $\hat{K}_T$  the set of all  $\tau \in \hat{K}$  such that  $[T, \tau] = 1$ . Then, as a representation of  $K$ ,  $(T|_K, \mathcal{H}_\tau)$  is equivalent with  $\tau$  for each  $\tau \in \hat{K}_T$ . We choose a complete orthonormal basis of  $\mathcal{H}$  such that

$$\{e_n^\tau; e_n^\tau \in \mathcal{H}_\tau, 1 \leq n \leq \dim \tau, \tau \in \hat{K}_T\}$$

and we denote by  $I$  the set of the indexes  $\{(\tau, n); 1 \leq n \leq \dim \tau, \tau \in \hat{K}_T\}$ . For each  $f \in \mathcal{H}$  the Fourier expansion of  $f$  is given by

$$f = \sum_{(\tau, n) \in I(f)} f_n^\tau e_n^\tau,$$

where  $f_n^\tau = \langle f, e_n^\tau \rangle_{\mathcal{H}}$  and  $I(f)$  the subset of  $I$  consisting of all  $(\tau, n)$  such that  $f_n^\tau \neq 0$ . Here we put

$$I_A(f) = \{(\tau, n); (T(\cdot)f)_n^\tau = \langle T(\cdot)f, e_n^\tau \rangle \text{ is not identically 0 on } A\}.$$

We say that  $\psi \in \mathcal{H}$  is admissible if there exist constants  $0 < C_1, C_2 < \infty$  such that, if  $(\tau, n) \in I_A(\psi)$ ,

$$C_1 \leq c_{\psi, \tau, n} = \int_A |\langle T(a)\psi, e_n^\tau \rangle|^2 D(a) da \leq C_2.$$

We put

$$\mathcal{H}_\psi = \{f \in \mathcal{H}; I(f) \subset I_A(\psi)\}.$$

Then, by using the bounded constants  $c_{\psi, \tau, n}$  we shall define a Fourier multiplier  $M_\psi$  on  $\mathcal{H}_\psi$  as follows. For each  $f = \sum_{(\tau, n) \in I(f)} f_n^\tau e_n^\tau$  in  $\mathcal{H}_\psi$

$$M_\psi f = \sum_{(\tau, n) \in I(f)} c_{\psi, \tau, n}^{-1/2} f_n^\tau e_n^\tau.$$

**Theorem 3.1.** Let  $\psi$  be admissible in  $\mathcal{H}$ . Then for any  $f \in \mathcal{H}_\psi$

$$(1) \quad C_1 \|f\|^2 \leq \int \int_{KA} |\langle f, T(ka)\psi \rangle|^2 D(a) dk da \leq C_2 \|f\|^2,$$

$$(2) \quad \|f\|^2 = \int \int_{KA} |\langle f, M_\psi T(ka)\psi \rangle|^2 D(a) dk da,$$

$$(3) \quad f = \int \int_{KA} \langle f, M_\psi T(ka)\psi \rangle M_\psi T(ka)\psi D(a) dk da.$$

*Proof.* We note that

$$T(k^{-1})f = \sum_{(\tau, n) \in I(f)} f_n^\tau T(k^{-1})e_n^\tau = \sum_{(\tau, n) \in I(f), (\tau', n') \in I} f_n^\tau \langle T(k^{-1})e_n^\tau, e_{n'}^{\tau'} \rangle e_{n'}^{\tau'}.$$

Then the orthogonality of the matrix coefficients of  $T|_K$  yields that

$$\begin{aligned} & \int \int_{KA} |\langle f, T(ka)\psi \rangle|^2 D(a) dk da \\ &= \int_A \sum_{(\tau, n) \in I(f)} |f_n^\tau|^2 |\langle T(a)\psi, e_n^\tau \rangle|^2 D(a) da \\ &= \sum_{(\tau, n) \in I(f)} |f_n^\tau|^2 \left( \int_A |\langle T(a)\psi, e_n^\tau \rangle|^2 D(a) da \right) \end{aligned}$$

Since

$$\|f\|^2 = \sum_{(\tau,n) \in I(f)} |f_n^\tau|^2 \quad \text{and} \quad I(f) \subset I_A(\psi),$$

(1) easily follows from the definition of the admissible vector  $\psi$ . We replace  $f$  by  $M_\psi f$  in the above calculation. Then  $|f_n^\tau|^2$  in the last equation turns to  $|f_n^\tau|^2 c_{\psi,\tau,n}^{-1}$  and then,  $c_{\psi,\tau,n}^{-1}$  cancels the integral over  $A$ . Thereby (2) follows. As for (3) we put  $\mathcal{H}(f) = \text{Span}\{e_n^\tau; (\tau, n) \in I(f)\}$  and define an operator  $Q$  on  $\mathcal{H}(f)$  by

$$h \mapsto \int \int_{KA} \langle f, M_\psi T(ka)\psi \rangle \langle h, M_\psi T(ka)\rangle \psi \, D(a) dk da.$$

Then (2) and the Schwarz inequality yield that  $Q$  is bounded and  $\|Q\| \leq \|f\|^2$ , and thereby, there exists  $f_0 \in \mathcal{H}(f)$  such that  $Q(h) = \langle h, f_0 \rangle$  and  $\|f_0\| = \|Q\|$ . Since  $Q(f) = \langle f, f_0 \rangle = \|f\|^2$  by (2), it easily follows that  $f = f_0$  (cf. [K]). Clearly,  $Q(h) = \langle h, f \rangle$  means (3).

**Remark 3.2.** When  $(T, \mathcal{H})$  is an irreducible square-integrable representation of  $G$ , it is well-known that each  $\psi \in \mathcal{H}$  is admissible and satisfies

$$c_{\psi,\tau,n} = d_T^{-1} \|\psi\|^2,$$

where  $c_T$  is the formal degree of  $T$  (cf. [V]). Furthermore, applying the orthogonality of the matrix coefficients on  $G$ , we can replace the integrals over  $KA$  in Theorem 3.1 by the ones over  $G$ .

§4. Example in  $SU(1, 1)$ .

Let  $G$  be  $SU(1, 1)$ . Then

$$K = \{k_\theta = \begin{pmatrix} e^{i\theta/2} & 0 \\ 0 & e^{-i\theta/2} \end{pmatrix}; 0 \leq \theta < 4\pi\},$$

$$A = \{a_t = \begin{pmatrix} \cosh t/2 & \sinh t/2 \\ \sinh t/2 & \cosh t/2 \end{pmatrix}; t \in \mathbf{R}\},$$

and  $A^+ = \{a_t; t > 0\}$ . In what follows we put

$$x = \tanh t.$$

Let  $(T_h, \mathcal{H}_h)$  ( $h \in \mathbf{Z}/2, h \geq 1$ ) be the holomorphic discrete series of  $G$  realized on the weighted Bergman space  $\mathcal{H}_h$  on the unit disk  $D = G/K$ :

$$\mathcal{H}_h = \{f : D \rightarrow \mathbf{C}; f \text{ is holomorphic on } D \text{ and}$$

$$\|f\|_h^2 = , (2h-1)^{-1} \int_D |f(z)|^2 (1-|z|^2)^{2(h-1)} dz < \infty\},$$

and  $(T_{1/2}, \mathcal{H}_{1/2})$  the limit of holomorphic discrete series of  $G$  realized on the Hardy space  $\mathcal{H}_{1/2}$  on  $D$ :

$$\mathcal{H}_{1/2} = \{f : D \rightarrow \mathbf{C}; f \text{ is holomorphic on } D \text{ and}$$

$$\|f\|_{1/2}^2 = \lim_{h \rightarrow 1/2} \|f\|_h^2 < \infty\}.$$

For  $h \in \mathbf{Z}/2, h \geq 1/2$  we denote by  $\langle \cdot, \cdot \rangle_h$  the inner product of  $\mathcal{H}_h$  and we put

$$e_n^h(z) = \left( \frac{, (2h+n)}{, (2h), (n+1)} \right)^{1/2} z^n \quad (n \in \mathbf{N}).$$

Then  $\{e_n^h; n \in \mathbf{N}\}$  is an orthonormal basis of  $\mathcal{H}_h$ . For simplicity we denote

$$\langle \cdot, \cdot \rangle = \langle \cdot, \cdot \rangle_{1/2} \text{ and } e_n(z) = e_n^{1/2}(z) = z^n.$$

According to this basis the matrix coefficients of  $T_h$  are given as follows (see [Sa]):

$$\begin{aligned} \langle T_h(g) e_n^h, e_m^h \rangle_h &= e^{i(n\theta+m\theta')} \langle T_h(a_t) e_n^h, e_m^h \rangle_h \quad (g = k_\theta a_t k_{\theta'}) \\ &= e^{i(n\theta+m\theta')} M(h; n, m; x), \end{aligned}$$

where for  $n \geq m$ ,

$$M(h; n, m; x) = C_{n,m}^h (1-x^2)^h (-x)^{n-m} F(-m, n+2h, n-m+1; x^2),$$

$$C_{n,m}^h = \left( \frac{, (n+1), (n+2h)}{, (m+1), (m+2h)} \right)^{1/2} \frac{1}{, (n-m+1)}$$

and  $F(a, b, c; x)$  is the hypergeometric function, and for  $m > n$  we change  $n$  and  $m$  by  $m$  and  $n$  respectively. Since

$$D(a_t) dt = \sinh(2t) dt = \frac{2x}{(1-x^2)^2} dx,$$

$M(h; n, m; x)$  ( $n, m \in \mathbf{N}$ ) are square-integrable on  $G$  if and only if  $h > 1/2$ . Here we note that for  $n \geq m$ ,

$$\begin{aligned} & \lim_{x \rightarrow 1} (1 - x^2)^{-h} M(h; n, m; x) \\ &= C_{n,m}^h (-1)^n \frac{(1 - m + n), (m + 2h)}{(2h), (n + 1)} \\ &= (-1)^n \frac{1}{(2h)} \left( \frac{(n + 2h), (m + 2h)}{(n + 1), (m + 1)} \right)^{1/2} \\ &= (-1)^n D_{n,m}^h \end{aligned}$$

and for  $m > n$ ,  $\lim_{x \rightarrow 1} (1 - x^2)^{-h} M(h; n, m; x) = (-1)^m D_{m,n}^h = (-1)^m D_{n,m}^h$ . Then we shall define the normalized matrix coefficients  $NM(h; n, m, x)$  as

$$NM(h; n, m; x) = (D_{n,m}^h)^{-1} M(h; n, m; x)$$

and the differences of the normalized matrix coefficients  $DM(h; n, m; x)$  as

$$DM(h; n, m; x) = NM(h; n, m; x) - NM(h; n + 2, m; x).$$

The key lemmas are the following.

**Lemma 4.1.** Let notations be as above. Then

$$\begin{aligned} DM(h; n, m; x) &= \frac{(1 - x^2)^{1/2}}{x} \\ &\times \left( \frac{m}{2h} NM(h + 1/2; n, m - 1; x) - \frac{m + 2h}{2h} NM(h + 1/2; n + 1, m; x) \right). \end{aligned}$$

*Proof.* We realize  $T_h$  on the circle and let  $z = e^{i\theta}$  ( $0 \leq \theta < 2\pi$ ) (see [Sa]). We first note that

$$\begin{aligned} (D_{n,m}^h)^{-1} e_n^h &= \left( \frac{(2h), (m + 1)}{(m + 2h)} \right)^{1/2} z^n, \\ (D_{n+2,m}^h)^{-1} e_{n+2}^h &= \left( \frac{(2h), (m + 1)}{(m + 2h)} \right)^{1/2} z^{n+2}, \end{aligned}$$

and moreover,

$$\begin{aligned}
& T_h(a_t)(z^n - z^{n+2}) \\
&= \frac{1}{(-z \sinh t/2 + \cosh t/2)^{2h}} \left( \frac{z \cosh t/2 - \sinh t/2}{-z \sinh t/2 + \cosh t/2} \right)^n \\
&\quad \times \left( 1 - \left( \frac{z \cosh t/2 - \sinh t/2}{-z \sinh t/2 + \cosh t/2} \right)^2 \right) \\
&= \frac{1}{(-z \sinh t/2 + \cosh t/2)^{2h}} \left( \frac{z \cosh t/2 - \sinh t/2}{-z \sinh t/2 + \cosh t/2} \right)^n \\
&\quad \times \frac{1 - z^2}{(-z \sinh t/2 + \cosh t/2)^2} \\
&= \frac{1}{\sinh t/2} \frac{1}{(-z \sinh t/2 + \cosh t/2)^{2h+1}} \left( \frac{z \cosh t/2 - \sinh t/2}{-z \sinh t/2 + \cosh t/2} \right)^n \\
&\quad \times \left( - \left( \frac{z \cosh t/2 - \sinh t/2}{-z \sinh t/2 + \cosh t/2} \right) + z \right).
\end{aligned}$$

On the other hand, we easily see that

$$\begin{aligned}
& \left\langle \left( \frac{, (2h), (m+1)}{, (m+2h)} \right)^{1/2} z^{n+1}, e_m^h \right\rangle_h \\
&= \langle (D_{n+1,m}^{h+1/2})^{-1} e_{n+1}^{h+1/2}, e_m^{h+1/2} \rangle_h \\
&= \frac{m+2h}{2h} \langle (D_{n+1,m}^{h+1/2})^{-1} e_{n+1}^{h+1/2}, e_m^{h+1/2} \rangle_{h+1/2}
\end{aligned}$$

and

$$\begin{aligned}
& \left\langle \left( \frac{, (2h), (m+1)}{, (m+2h)} \right)^{1/2} z^n, e_{m-1}^h \right\rangle_h \\
&= \langle (D_{n,m-1}^{h+1/2})^{-1} e_n^{h+1/2}, e_{m-1}^{h+1/2} \rangle_h \\
&= \frac{m}{2h} \langle (D_{n,m-1}^{h+1/2})^{-1} e_n^{h+1/2}, e_{m-1}^{h+1/2} \rangle_{h+1/2}.
\end{aligned}$$

Then the desired result follows.

**Lemma 4.2.** Let notations be as above. Then for each  $n, m \in \mathbf{N}$ ,

$$0 < \int_0^1 DM(h; n, m; x)^2 \frac{2x}{(1-x^2)^2} dx < \infty,$$

and especially, for  $m > n$

$$\begin{aligned} & \int_0^1 DM(h; n, m; x)^2 \frac{2x}{(1-x^2)^2} dx \\ &= , (2h)^2 2(n+h+1) \frac{(m+1)}{(m+2h)} \frac{(n+1)}{(n+2h+2)}. \end{aligned}$$

*Proof.* The case of  $m > n$ : We note that

$$\begin{aligned} & \frac{(1-x^2)^{1/2}}{x} \frac{m}{2h} NM(h+1/2; n, m-1; x) \\ &= Ax^{m-n-2} (1-x^2)^{h+1} G_n(m-n+2h, m-n; x^2) \end{aligned}$$

and

$$\begin{aligned} & \frac{(1-x^2)^{1/2}}{x} \frac{m+2h}{2h} NM(h+1/2; n+1, m; x) \\ &= \frac{m+2h}{n+2h+1} Ax^{m-n-2} (1-x^2)^{h+1} G_{n+1}(m-n+2h, m-n; x^2), \end{aligned}$$

where

$$A = \frac{, (2h), (m+1)}{, (n+2h+1), (m-n)}$$

and  $G_n(x) = G_n(\alpha, \gamma, x)$  ( $\alpha = m-n+2h, \gamma = m-n$ ) is the Jacobi polynomial. Hence,

$$\begin{aligned} I &= \int_0^1 DM(h; n, m; x)^2 \frac{2x}{(1-x^2)^2} dx \\ &= A^2 \int_0^1 x^{2(m-n-2)} (1-x^2)^{2h} \left( G_n(x^2) - \frac{m+2h}{n+2h+1} G_{n+1}(x^2) \right)^2 2x dx \\ &= A^2 \int_0^1 x^{\gamma-1} (1-x)^{\alpha-\gamma} \left( G_n(x) - \frac{m+2h}{n+2h+1} G_{n+1}(x) \right)^2 \frac{dx}{x}. \end{aligned}$$

We here consider the case of  $m > n+1$ . Then,  $\gamma-2 = m-n-2 \geq 0$ . We note that  $G_n^2 = (G_n - 1)G_n + G_n$  and  $(G_n - 1)/x$  is the polynomial of

degree  $n - 1$ . So the orthogonality relations for the Jacobi polynomials and the definition of  $G_n(x)$ ;

$$G_n(x) = \frac{, (\gamma)}{, (\gamma + n)} x^{1-\gamma} (1-x)^{\gamma-\alpha} \left( \frac{d}{dx} \right)^n \left( x^{\gamma+n-1} (1-x)^{\alpha+n-\gamma} \right)$$

yield that

$$\begin{aligned} & \int_0^1 x^{\gamma-1} (1-x)^{\alpha-\gamma} G_n(x)^2 \frac{dx}{x} \\ &= \int_0^1 x^{\gamma-1} (1-x)^{\alpha-\gamma} G_n(x) \frac{dx}{x} \\ &= , (m-n)^2 \frac{, (n+1), (n+2h+1)}{, (m+1), (m+2h)} \frac{m}{m-n-1} \\ &= B, \end{aligned}$$

and similarly,

$$\begin{aligned} & \int_0^1 x^{\gamma-1} (1-x)^{\alpha-\gamma} G_{n+1}(x)^2 \frac{dx}{x} \\ &= \int_0^1 x^{\gamma-1} (1-x)^{\alpha-\gamma} G_n(x) G_{n+1}(x) \frac{dx}{x} \\ &= \int_0^1 x^{\gamma-1} (1-x)^{\alpha-\gamma} G_{n+1}(x) \frac{dx}{x} \\ &= \frac{n+2h+1}{m+2h} \frac{n+1}{m} B. \end{aligned}$$

Therefore,

$$\begin{aligned} I &= A^2 B \left( 1 - 2 \frac{n+1}{m} + \frac{n+1}{m} \frac{m+2h}{n+2h+1} \right) \\ &= A^2 B \frac{2(m-n-1)(n+h+1)}{m(n+2h+1)} \end{aligned}$$

and hence, the desired result follows.

In the case of  $m = n+1$  we note that  $(G_n(x) - G_{n+1}(x))/x$  is a polynomial of degree  $n$  and thus, the integral  $I$  is well-defined. Then the analytic continuation on  $\gamma$ , letting  $\gamma \rightarrow 1$  in the previous case, yields the desired formula for  $m = n+1$ .

The case of  $m \leq n$ : Since  $M(h + 1/2; n, m - 1; x)$  and  $M(h + 1/2; n + 1, m; x)$  have the term  $x^{n-m+1}$  and  $n - m + 1 \geq 1$ , it easily follows from Lemma 4.1 that the desired integral is positive and finite.

This completes the proof of the lemma.

**Lemma 4.3.** Let notations be as above and suppose that

$$n, m \in 2\mathbf{N} \quad \text{or} \quad n, m \in 2\mathbf{N} + 1.$$

Then, for  $p > n, m$

$$\begin{aligned} & \int_0^1 DM(h; n, p, x) DM(h; m, p, x) \frac{2x}{(1-x^2)^2} dx \\ &= \delta_{nm}, (2h)^2 2(n+h+1) \frac{, (p+1)}{, (p+2h)} \frac{, (n+1)}{, (n+2h+2)}. \end{aligned}$$

*Proof.* When  $n = m$ , it follows from Lemma 4.2. We may suppose that  $n > m$  and hence,  $n - m \geq 2$  and even. Then, applying the same argument used in the proof of Lemma 4.2, we see that the desired integral equals to

$$\begin{aligned} & \int_0^1 x^{p-n-1+(n-m)/2} (1-x)^{2h} \\ & \times \left( G_n(x) - \frac{p+2h}{n+2h+1} G_{n+1}(x) \right) \left( G_m(x) - \frac{p+2h}{m+2h+1} G_{m+1}(x) \right) \frac{dx}{x}. \end{aligned}$$

Since  $(n - m)/2$  is integer,  $0 \leq (n - m)/2 - 1 \leq n - 1$ , and

$$\left( G_m(x) - \frac{p+2h}{m+2h+1} G_{m+1}(x) \right)$$

is a polynomial of degree  $m+1 < n$ , the orthogonality relations for the Jacobi polynomials yield that the integral equals to 0.

We here note that, if  $h = 1/2$ , then  $D_{n,m}^h = 1$  and hence,

$$\begin{aligned} DM(1/2; n, m; x) &= M(1/2; n, m; x) - M(1/2; n + 2, m; x) \\ &= \langle T_{1/2}(a_t)(e_n - e_{n+2}), e_m \rangle. \end{aligned}$$

Therefore, Lemma 4.2 implies that

$$0 < \int_A |\langle T_{1/2}(a_t)(e_n - e_{n+2}), e_m \rangle|^2 D(a_t) dt < \infty$$

and for  $m > n$  this integral equals to

$$\frac{(2n+3)}{(n+1)(n+2)}.$$

Furthermore, these differences  $\langle T_{1/2}(a_t)(e_n - e_{n+2}), e_m \rangle$  satisfy the quasi-orthogonality relations stated in Lemma 4.3 with  $h = 1/2$ . Thereby, as an application of Theorem 3.1, we see the following.

**Theorem 4.4.** Let  $G = SU(1, 1)$  and  $(T_{1/2}, \mathcal{H}_{1/2})$  the limit of the discrete series of  $G$ .

(1) Let  $\psi$  be a finite linear combination of  $e_{n+2} - e_n$ . Then there exist constants  $0 < C_1, C_2 < \infty$  such that for any  $f$  in  $\mathcal{H}_{1/2}$

$$C_1 \|f\|^2 \leq \int \int_{KA} |\langle f, T_{1/2}(ka_t)\psi \rangle|^2 \sinh 2t \ dk dt \leq C_2 \|f\|^2.$$

(2) Let

$$\psi = \sum c_n \left( \frac{(2n+3)}{(n+1)(n+2)} \right)^{-1/2} (e_{n+2} - e_n),$$

where the sum is taken over  $0 \leq n \leq N, n \in 2\mathbb{N}$  or  $0 \leq n \leq N, n \in 2\mathbb{N} + 1$ , and let  $\|\psi\|_0^2 = \sum |c_n|^2$ . Then for any  $f$  in the  $L^2$ -span of  $\{e_p, p \geq N + 1\}$ ,

$$f(x) = \frac{1}{\|\psi\|_0} \int \int_{KA} \langle f, T_{1/2}(ka_t)\psi \rangle T_{1/2}(ka_t)\psi \sinh 2t \ dk dt.$$

## References

[AAG] S. T. Ali, J.-P. Antoine and J.-P. Gazeau, *Square integrability of group representations on homogeneous spaces I, II*, Ann. Inst. H. Poincaré, Vol. 55, 1991, pp. 829-855, *ibid.*, Ann. Inst. H. Poincaré, Vol. 55, 1991, pp. 857-890.

- [Ba] V. Bargmann, *On unitary ray representations for continuous groups*, Ann. Math., Vol. 48, 1947, pp. 568-640.
- [BT] D. Bernier and K. F. Taylor, *Wavelets from square-integrable representations*, SIAM J. Math. Anal., Vol. 27, 1996, pp. 594-608.
- [Bo] G. Bohnke, *Treillis d'ondelettes associés aux groupes de Lorentz*, Ann. Inst. H. Poincaré, Vol. 54, 1991, pp. 245-259.
- [DB] De Bièvre, *coherent states over symplectic homogeneous spaces*, J. Math. Phys., Vol. 7, 1989, pp. 1401-1407.
- [DM] G. B. Duflo and C.C. Moore, *On the regular representation of a nonunimodular locally compact group*, J. Funct. Anal., Vol. 21, 1976, pp. 209-243.
- [G] R. Godement, *Sur les relations d'orthogonalité de Bargmann*, C. R. Acad. Sci., Vol. 255, 1947, pp. 521-523, *ibid.*, pp. 657-659.
- [GMP] A. Grossmann, J. Morlet, and T. Paul, *Transforms associated to square integrable group representations. I*, J. Math. Phys., Vol. 26, 1985, pp. 2473-2479.
- [HC] Harish-Chandra, *Discrete series for semisimple Lie groups I, II*, Acta Math, Vol. 113, 1965, pp. 241-318, *ibid.*, Vol. 116, 1966, pp. 1-111.
- [K1] T. Kawazoe, *Wavelet transforms associated to a principal series representation of semisimple Lie groups I, II*, Proc. Japan Acad. Vol. 71, 1995, pp. 154-157, *ibid.*, Proc. Japan Acad. Vol. 71, 1995, pp. 158-160.
- [K2] T. Kawazoe, *Wavelet transforms associated to an induced representation of  $SL(n+2, R)$* , Ann. Inst. H. Poincaré, Vol. 65, 1996, pp. 1-13.
- [K3] T. Kawazoe, *Wavelet transforms associated to the analytic continuation of the holomorphic discrete series of a semisimple Lie group*, Keio Research Report, 1998.
- [KT] C. Kalisa and B. Torrésani,  *$N$ -dimensional affine Weyl-Heisenberg wavelets*, Ann. Inst. H. Poincaré, Vol. 59, 1993, pp. 201-236.

- [L] H. Liu, *Wavelet transforms and symmetric tube domains*, J. Lie Theory, Vol. 8, 1998, pp. 351-366.
- [M] R. Murenzi, *Ondelettes multidimensionnelles et application à l'analyse d'images*, Ph. D. Thesis, Louvain-la-Neuve, 1990.
- [MW] C. C. Moore and J. A. Wolf, *Square integrable representations on nilpotent groups*, Trans. Amer. Math. Soc., Vol. 185, 1973, pp. 445-462.
- [Sa] P. J. Sally, *Analytic Continuation of the Irreducible Unitary Representations of the Universal Covering Group of  $SL(2, \mathbf{R})$* , Memoirs of A. M. S., Vol. 69, American Mathematical Society, Providence, Rhode Island, 1967.
- [Su] M. Sugiura, *Unitary Representations and Harmonic Analysis, An Introduction*, Second Edition, North-Holland/Kodansha, Amsterdam/Tokyo, 1990.
- [T1] B. Torrésani, *Wavelets associated with representations of the affine Weyl-Heisenberg group*, J. Math. Phys., Vol. 32, 1991, pp. 1273-1279.
- [T2] B. Torrésani, *Time-frequency representation: wavelet packets and optimal decomposition*, Ann. Inst. H. Poincaré, Vol. 56, 1992, pp. 215-234.
- [U] A. Unterberger and J. Unterberger, *A quantization of the Cartan Domain BDI ( $q = 2$ ) and operators on the light cone*, J. Funct. Anal., Vol. 72, 1987, pp. 279-319.
- [V] V. S. Varadarajan, *Harmonic Analysis on Real Reductive Groups*, Lecture Note in Math., Springer, Berlin, New York, Vol. 576, 1977.
- [WZ] Y. Wu and X. Zhong, *Discrete wavelet transform on circles*, preprint.
- [AV1] J-P. Antoine and P. Vandergheynst, *Wavelets on the  $n$ -sphere and other manifolds*, J. Math. Phys., Vol. 39, 1998, pp. 3987-4008.
- [AV2] J-P. Antoine and P. Vandergheynst, *Wavelets on the 2-sphere: A group-theoretical approach*, Appl. Comput. Harmon. Anal., Vol. 7, 1999, pp. 262-291.